


BOTOX/DYSPORT

Botulinum Toxin A



Botox/dysport are comprised of botulinum toxin A, which relaxes muscles in the specific area that is injected. Both work by decreasing the force of muscle contraction in specific areas of your face.

Areas That Can Be Treated With BOTOX/DYSPORT

- ~ Forehead lines
- ~ Crows feet and fine lines around the eyes
- ~ Frown lines between the eyes or "11" furrows
- ~ Droopy eyebrows
- ~ Marionette area and wrinkles around the mouth/upper lip
- ~ Chin dimpling
- ~ Platysmal bands or neck bands commonly known as Nefertiti Lift

Botox/Dysport are great for those who struggles with the signs of aging and also can be used as preventative measures due to genetic predisposition for severe wrinkling. These injectables can give you a younger-looking appearance with a smoother forehead area diminishing lines and wrinkles without the for fsurgery or recovery. You will begin to see improvement in the treated area in 48 hours, with full effects at about two weeks. Say goodbye to your wrinkles for 3-5 months!

Cost:

- \$12 per unit

THESE INJECTABLES CAN GIVE YOU A YOUNGER LOOKING APPEARANCE WITH A SMOOTHER FOREHEAD AREA DIMINISHING LINES AND WRINKLES WITHOUT RECOVERY.

PRE-TREATMENT INSTRUCTIONS

- Avoid sun exposure prior to treatment. Sunburned skin cannot be treated.
- Do not bleach, wax, tweeze or use depilatory creams near the area for 1 to 2 days prior to treatment.
- Discontinue use of retinoids (Retin A, Tazorac, Differin, etc.) 1 to 2 days prior to treatment.
- Discontinue use of all 10% or higher alpha hydroxy acids, exfoliants, etc. for 1 to 2 days prior to treatment.

- If you have a history of perioral herpes, begin prophylactic antiviral therapy of the day before treatment.

DAY OF TREATMENT

- Omit lotions, creams, makeup or deodorant around the area to be treated.
- Inform your provider any changes in medical history and of all medications you're taking.

POST TREATMENT

- Redness might be present (and last several hours) immediately after treatment.
- Bruising, swelling and/or redness may occur.
- Avoid sun exposure after the treatment to reduce the chance of hyperpigmentation.
- Remain upright for at least four hours post treatment. No heavy lifting, bending over, exercising, massage, head bands or glasses pressing against areas of injection. Since neuromodulators can take up to two weeks for completion it is recommended no facial treatments or facial massages be performed within that two-week period.
- Avoid blood thinners after treatment to reduce the chance of bruising. Tylenol is acceptable.
- If for any reason, scabbing should occur, do not pick or scratch at the treated area. Treat skin gently. Wash with a gentle cleanser, cool water, using hands only. Pat dry.
- Follow RN instructions on caring for skin post treatment.
- Use sunscreen on a daily basis. This is extremely important!

